




## Identifying Specific Bacteria Group

While the common V3-V4 target region is well-sequenced, the identification of microbial community is still not fully understood.

The following table shows a compilation of recommended 16s rRNA region for identifying specific bacteria group in different area of research.



Research Area	16s rRNA region	Targeted Bacteria Group	Citation
Pathogenic Bacteria	V1	<i>Staphylococcus aureus</i> , coagulase negative <i>Staphylococcus</i> sp.	Chakravorty, Helb et al. 2007
	V2	<i>Mycobacterium</i> species	Chakravorty, Helb et al. 2007
	V3	<i>Haemophilus</i> species	Chakravorty, Helb et al. 2007
Subgingival Microbial Communities	V1-V3	Genera <i>Prevotella</i> , <i>Fusobacterium</i> , <i>Streptococcus</i> , <i>Granulicatella</i> , <i>Bacteroides</i> , <i>Porphyromonas</i> and <i>Treponema</i>	Kumar, Brooker et al. 2011
	V7-V9	Genera <i>Veillonella</i> , <i>Streptococcus</i> , <i>Eubacterium</i> , <i>Enterococcus</i> , <i>Treponema</i> , <i>Catonella</i> and <i>Selenomonas</i>	Kumar, Brooker et al. 2011
Paddy Soil Microbiome	V1-V3	Actinobacteria (particularly in genus <i>Arthrobacter</i> )	Li, Zhang et al. 2009
	V3	$\beta$ -Proteobacteria, (particularly in genus <i>Gallionella</i> ) and $\alpha$ -Proteobacteria	Li, Zhang et al. 2009
	V3-V5	Actinobacteria (particularly in genus <i>Arthrobacter</i> )	Li, Zhang et al. 2009
	V8	$\beta$ -Proteobacteria, $\gamma$ -Proteobacteria and particularly in genus <i>Acinetobacter</i>	Li, Zhang et al. 2009
	V6-V8	Chlamydiae and $\beta$ -Proteobacteria	Li, Zhang et al. 2009

## Conclusion

Depending on the research area as shown in the papers reviewed above, the other 16s rRNA regions that were sequenced resulted in different matches of bacteria species in the database. With V3-V4 region being widely sequenced and its results well-established, it is highly recommended to sequence more hypervariable regions in combination with this region for a more complete microbial community profile.

In addition, the price of next generation sequencing has decreased over the years, thus making it more affordable for researchers to do so now, more than ever.

Hence, for an improved overall representation of the bacterial community, sequencing more individual or grouped 16s rRNA regions will allow researchers to achieve that.

### Reference

- Chakravorty, S., Helb, D., Burday, M., Connell, N., & Alland, D. (2007). A detailed analysis of 16S ribosomal RNA gene segments for the diagnosis of pathogenic bacteria. *Journal of Microbiological Methods*, 69(2), 330-339. doi:10.1016/j.mimet.2007.02.005
- Kumar, P. S., et al. (2011). "Target region selection is a critical determinant of community fingerprints generated by 16S pyrosequencing." *PLoS One* 6(6): e20956.
- Zheng, W., Tsompana, M., Ruscitto, A., Sharma, A., Genco, R., Sun, Y., & Buck, M. J. (2015). An accurate and efficient experimental approach for characterization of the complex oral microbiota. *Microbiome*, 3(1). doi:10.1186/s40168-015-0110-9
- Li, H., Zhang, Y., Li, D., Xu, H., Chen, G., & Zhang, C. (2009). Comparisons of different hypervariable regions of *rrs* genes for fingerprinting of microbial communities in paddy soils. *Soil Biology and Biochemistry*, 41(5), 954-968. doi:10.1016/j.soilbio.2008.10.030
- Hong, S., Bunge, J., Leslin, C., Jeon, S., & Epstein, S. S. (2009). Polymerase chain reaction primers miss half of rRNA microbial diversity. *The ISME Journal*, 3(12), 1365-1373. doi:10.1038/ismej.2009.89
- Sipos, R., Székely, A. J., Palatinszky, M., Révész, S., Márialigeti, K., & Nikolausz, M. (2007). Effect of primer mismatch, annealing temperature and PCR cycle number on 16S rRNA gene-targeting bacterial community analysis. *FEMS Microbiology Ecology*, 60(2), 341-350. doi:10.1111/j.1574-6941.2007.00283.x
- Assessment of biases associated with profiling simple, model communities using terminal-restriction fragment length polymorphism-based analyses. *Journal of Microbiological Methods*, 67(1), 9-19. doi:10.1016/j.mimet.2006.02.011